

## PHENTOLAMINE MESYLATE - phentolamine mesylate injection, powder, for solution


Bedford Laboratories

### Rx ONLY.

### DESCRIPTION

Phentolamine Mesylate for Injection USP, is an antihypertensive, available in vials for intravenous and intramuscular administration. Each vial contains phentolamine mesylate USP, 5 mg and mannitol USP, 25 mg in sterile, lyophilized form.

Phentolamine mesylate is *m*-[*N*-(2-Imidazolin-2-ylmethyl)-*p*-toluidino]phenol monomethanesulfonate (salt), and its structural formula is:



Molecular Formula - C<sub>17</sub>H<sub>19</sub>N<sub>3</sub>O•CH<sub>4</sub>O<sub>3</sub>S

M.W. - 377.47

Phentolamine mesylate USP is a white or off-white, odorless crystalline powder. Its solutions are acid to litmus. It is freely soluble in water and in alcohol, and slightly soluble in chloroform. It melts at about 178°C.

### CLINICAL PHARMACOLOGY

Phentolamine mesylate produces an alpha-adrenergic block of relatively short duration. It also has direct, but less marked, positive inotropic and chronotropic effects on cardiac muscle and vasodilator effects on vascular smooth muscle.

Phentolamine has a half-life in the blood of 19 minutes following intravenous administration. Approximately 13% of a single intravenous dose appears in the urine as unchanged drug.

### INDICATIONS AND USAGE

Phentolamine Mesylate for Injection is indicated for the prevention or control of hypertensive episodes that may occur in a patient with pheochromocytoma as a result of stress or manipulation during preoperative preparation and surgical excision.

Phentolamine Mesylate for Injection is indicated for the prevention or treatment of dermal necrosis and sloughing following intravenous administration or extravasation of norepinephrine.

Phentolamine Mesylate for Injection is also indicated for the diagnosis of pheochromocytoma by the phentolamine blocking test.

### CONTRAINDICATIONS

Myocardial infarction, history of myocardial infarction, coronary insufficiency, angina, or other evidence suggestive of coronary artery disease; hypersensitivity to phentolamine or related compounds.

### WARNINGS

Myocardial infarction, cerebrovascular spasm, and cerebrovascular occlusion have been reported to occur following the administration of phentolamine, usually in association with marked hypotensive episodes.

For screening tests in patients with hypertension, the generally available urinary assay of catecholamines or other biochemical assays have largely replaced the phentolamine and other pharmacological tests for reasons of accuracy and safety. None of the chemical or pharmacological tests is infallible in the diagnosis of pheochromocytoma. The phentolamine blocking test is not the procedure of choice and should be reserved for cases in which additional confirmatory evidence is necessary and the relative risks involved in conducting the test have been considered.

### PRECAUTIONS

#### General

Tachycardia and cardiac arrhythmias may occur with the use of phentolamine or other alpha-adrenergic blocking agents. When possible, administration of cardiac glycosides should be deferred until cardiac rhythm returns to normal.

#### Drug Interactions

See DOSAGE AND ADMINISTRATION. Diagnosis of pheochromocytoma, Preparation.

#### Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term carcinogenicity studies, mutagenicity studies, and fertility studies have not been conducted with phentolamine.

## Pregnancy

### Teratogenic Effects-Pregnancy Category C

Administration of phentolamine to pregnant rats and mice at oral doses 24 to 30 times the usual daily human dose (based on a 60 kg human) resulted in slightly decreased growth and slight skeletal immaturity of the fetuses. Immaturity was manifested by increased incidence of incomplete or unossified calcanei and phalangeal nuclei of the hind limb and of incompletely ossified sternbrae. At oral doses 60 times the usual daily human dose (based on a 60 kg human), a slightly lower rate of implantation was found in the rat. Phentolamine did not affect embryonic or fetal development in the rabbit at oral doses 20 times the usual daily human dose (based on a 60 kg human). No teratogenic or embryotoxic effects were observed in the rat, mouse, or rabbit studies.

There are no adequate and well-controlled studies in pregnant women. Phentolamine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from phentolamine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

## Pediatric Use

See DOSAGE AND ADMINISTRATION.

## ADVERSE REACTIONS

Acute and prolonged hypotensive episodes, tachycardia, and cardiac arrhythmias have been reported. In addition, weakness, dizziness, flushing, orthostatic hypotension, nasal stuffiness, nausea, vomiting, and diarrhea may occur.

## OVERDOSAGE

### Acute Toxicity

No deaths due to acute poisoning with phentolamine have been reported.

Oral LD<sub>50</sub>'s (mg/kg): mice, 1000; rats, 1250.

## Signs and Symptoms

Overdosage with phentolamine is characterized chiefly by cardiovascular disturbances, such as arrhythmias, tachycardia, hypotension, and possibly shock. In addition, the following might occur: excitation, headache, sweating, pupillary contraction, visual disturbances; nausea, vomiting, diarrhea; hypoglycemia.

## Treatment

There is no specific antidote.

A decrease in blood pressure to dangerous levels or other evidence of shocklike conditions should be treated vigorously and promptly. The patient's legs should be kept raised and a plasma expander should be administered. If necessary, intravenous infusion of norepinephrine, titrated to maintain blood pressure at the normotensive level, and all available supportive measures should be included. Epinephrine should not be used, since it may cause a paradoxical reduction in blood pressure.

## DOSAGE AND ADMINISTRATION

*The reconstituted solution should be used upon preparation and should not be stored.*

- 1. Prevention or control of hypertensive episodes in the patient with pheochromocytoma.** For preoperative reduction of elevated blood pressure, 5 mg of phentolamine mesylate (1 mg for children) is injected intravenously or intramuscularly 1 or 2 hours before surgery, and repeated if necessary.  
During surgery, phentolamine mesylate (5 mg for adults, 1 mg for children) is administered intravenously as indicated, to help prevent or control paroxysms of hypertension, tachycardia, respiratory depression, convulsions, or other effects of epinephrine intoxication. (Postoperatively, norepinephrine may be given to control the hypotension that commonly follows complete removal of a pheochromocytoma.)
- 2. Prevention or treatment of dermal necrosis and sloughing following intravenous administration or extravasation of norepinephrine.**  
*For Prevention:* 10 mg of phentolamine mesylate is added to each liter of solution containing norepinephrine. The pressor effect of norepinephrine is not affected.  
*For Treatment:* 5 to 10 mg of phentolamine mesylate in 10 mL of saline is injected into the area of extravasation within 12 hours.
- 3. Diagnosis of pheochromocytoma - phentolamine blocking test.**

The test is most reliable in detecting pheochromocytoma in patients with sustained hypertension and least reliable in those with paroxysmal hypertension. False-positive tests may occur in patients with hypertension without pheochromocytoma.

#### a. Intravenous

##### Preparation

The CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS sections should be reviewed. Sedatives, analgesics, and all other medications except those that might be deemed essential (such as digitalis and insulin) are withheld for at least 24 hours, and preferably 48 to 72 hours, prior to the test. Antihypertensive drugs are withheld until blood pressure returns to the untreated, hypertensive level. This test is not preformed on a patient who is normotensive.

##### Procedure

The patient is kept at rest in a supine position throughout the test, preferably in a quiet, darkened room. Injection of phentolamine is delayed until blood pressure is stabilized, as evidenced by blood pressure readings taken every 10 minutes for at least 30 minutes.

Five milligrams of phentolamine mesylate is dissolved in 1 mL of Sterile Water for Injection. The dose for adults is 5 mg; for children, 1 mg.

The syringe needle is inserted into the vein, and injection is delayed until pressor response to venipuncture has subsided.

Phentolamine is injected rapidly. Blood pressure is recorded immediately after injection, at 30-second intervals for the first 3 minutes, and at 60-second intervals for the next 7 minutes.

##### Interpretation

A positive response, suggestive of pheochromocytoma, is indicated when the blood pressure is reduced more than 35 mm Hg systolic and 25 mm Hg diastolic. A typical positive response is a reduction in pressure of 60 mm Hg systolic and 25 mm Hg diastolic. Usually, maximal effect is evident within 2 minutes after injection. A return to preinjection pressure commonly occurs within 15 to 30 minutes but may occur more rapidly.

If blood pressure decreases to a dangerous level, the patient should be treated as outlined under OVERDOSAGE.

A positive response should always be confirmed by other diagnostic procedures, preferably by measurement of urinary catecholamines or their metabolites.

A negative response is indicated when the blood pressure is elevated, unchanged, or reduced less than 35 mm Hg systolic and 25 mm Hg diastolic after injection of phentolamine. A negative response to this test does not exclude the diagnosis of pheo-chromocytoma, especially in patients with paroxysmal hypertension in whom the incidence of false-negative responses is high.

#### b. Intramuscular

If the intramuscular test for pheochromocytoma is preferred, preparation is the same as for the intravenous test. Five milligrams of phentolamine mesylate is then dissolved in 1 mL of Sterile Water for Injection. The dose for adults is 5 mg intramuscularly; for children, 3 mg. Blood pressure is recorded every 5 minutes for 30 to 45 minutes following injection. A positive response is indicated when the blood pressure is reduced 35 mm Hg systolic and 25 mm Hg diastolic, or more, within 20 minutes following injection.

*Note:* Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

#### HOW SUPPLIED

Phentolamine Mesylate for Injection USP, 5 mg, for intramuscular or intravenous use, is supplied in a 2mL vial and individually boxed. **NDC 55390-113-01.**

The reconstituted solution should be used upon preparation and should not be stored.

Store at controlled room temperature, 15° to 30°C (59° to 86°F).

Manufactured for:

Bedford Laboratories™

Bedford, OH 44146

May 1999


Manufactured by:

Ben Venue Laboratories™

Bedford, OH 44146


PHT-P01

#### VIAL LABEL



Vial Label 5 mg

# CARTON



Unit Carton 5 mg